

# Interleaving operator $\|\|$ for TS

PC2.2-DEF-INTERLEAVING-TS

$$\mathcal{T}_1 = (S_1, Act_1, \longrightarrow_1, S_{0,1}, AP_1, L_1)$$

$$\mathcal{T}_2 = (S_2, Act_2, \longrightarrow_2, S_{0,2}, AP_2, L_2)$$

The transition system  $\mathcal{T}_1 \|\| \mathcal{T}_2$  is defined by:

$$\mathcal{T}_1 \|\| \mathcal{T}_2 = (S_1 \times S_2, Act_1 \cup Act_2, \longrightarrow, S_{0,1} \times S_{0,2}, AP, L)$$

where the transition relation  $\longrightarrow$  is given by:

$$\frac{s_1 \xrightarrow{\alpha} s'_1}{\langle s_1, s_2 \rangle \xrightarrow{\alpha} \langle s'_1, s_2 \rangle} \quad \frac{s_2 \xrightarrow{\alpha} s'_2}{\langle s_1, s_2 \rangle \xrightarrow{\alpha} \langle s_1, s'_2 \rangle}$$

atomic propositions:  $AP = AP_1 \uplus AP_2$

labeling function:  $L(\langle s_1, s_2 \rangle) = L_1(s_1) \cup L_2(s_2)$